

SUPPLEMENTARY.

Table S1. Latencies of the P3 component in mentally healthy individuals (control group) and schizophrenia patients

Group	Condition	Electrode								
		F3	F4	Fz	C3	C4	Cz	P3	P4	Pz
C	50% S	280.1 (44.0)	290.0 (46.1)	286.5 (44.5)	305.3 (47.6)	302.6 (48.1)	298.9 (46.6)	321.4 (62.9)	310.8 (50.7)	321.1 (59.5)
SZ		279.8 (46.5)	287.9 (53.7)	283.5 (47.0)	278.7 (43.6)	285.7 (44.6)	281.2 (46.8)	283.8 (42.2)	285.3 (43.3)	283.9 (41.9)
C	50% D	286.8 (39.5)	286.5 (40.6.)	285.4 (41.8)	290.4 (36.7)	289.3 (38.9)	295.3 (41.6)	300.7 (39.4)	298.3 (38.9)	306.1 (40.7)
SZ		270.8 (43.2)	273.4 (39.9)	282.3 (44.9)	273.1 (37.4)	278.5 (37.2)	277.9 (39.3)	288.4 (45.2)	291.2 (35.0)	289.8 (36.5)
C	80% S	287.6 (57.9)	284.7 (53.5)	287.9 (55.6)	301.4 (53.2)	302.9 (55.0)	309.6 (53.0)	309.9 (55.3)	307.7 (52.1)	319.8 (50.0)
SZ		280.2 (49.6)	283.8 (48.6)	282.9 (52.0)	281.4 (43.9)	281.0 (40.6)	284.0 (46.4)	285.3 (38.1)	292.0 (35.7)	296.9 (43.6)
C	80% D	285.9 (36.5)	290.9 (36.0)	284.5 (34.2)	290.3 (35.5)	300.0 (33.3)	294.5 (32.0)	308.5 (44.8)	313.3 (41.5)	310.7 (45.1)
SZ		278.8 (41.2)	280.2 (41.9)	282.6 (44.3)	278.2 (42.5)	287.3 (41.6)	281.0 (42.7)	294.9 (55.1)	299.5 (39.3)	296.5 (40.2)

Note: The description takes into account the arithmetic mean (standard deviation). C — control group (*n*=20), D — deviant stimulus, S — standard stimulus, SZ — patients with schizophrenia (*n*=18).