

SUPPLEMENTARY.

Table S2. Comparative characteristics of clusters by all study parameters

Variable, statistics	Response to categorical variable	Cluster 1 (n=41)	Cluster 2 (n=39)	Cluster 3 (n=27)	Cluster 4 (n=8)	Total (n=115)	Test
Min of BMI (Median [Q1; Q3])	—	13.90 [12.60; 15.65]	15.40 [14.10; 16.70]	14.00 [11.65; 16.45]	14.50 [13.62; 15.75]	14.57 [12.65; 16.30]	Kruskal: H=7.429, p=0.059, $\epsilon^2=0.065$
Age (Median [Q1; Q3])	—	21.00 [18.00; 24.00]	22.00 [19.25; 26.00]	27.00 [21.00; 31.00]	21.50 [20.75; 29.50]	22.00 [19.00; 26.00]	Kruskal: H=9.482, p=0.024, $\epsilon^2=0.086$
BMI (Mean [SD])	—	14.16 (2.02)	15.45 (1.91)	14.14 (2.68)	14.71 (2.13)	14.63 (2.21)	ANOVA: F=2.825, p=0.042, d=0.593
Duration of the disease, years (Median [Q1; Q3])	—	4.00 [1.00; 7.50]	5.00 [3.00; 9.00]	6.00 [3.00; 9.00]	3.50 [1.75; 5.50]	5.00 [2.00; 9.00]	Kruskal: H=5.459, p=0.141, $\epsilon^2=0.050$
Checklist of symptoms (n=115)							
Dysphoria, n (%)	0	37 (90.2%)	16 (41.0%)	3 (11.1%)	5 (62.5%)	61 (100.0%)	Chi ² =44.389, p=0.000, V=0.621
	1	4 (9.8%)	23 (59.0%)	24 (88.9%)	3 (37.5%)	54 (100.0%)	
Mood swings, n (%)	0	26 (63.4%)	26 (66.7%)	12 (44.4%)	7 (87.5%)	71 (100.0%)	Chi ² =6.116, p=0.106, V=0.231
	1	15 (36.6%)	13 (33.3%)	15 (55.6%)	1 (12.5%)	44 (100.0%)	
Anxiety, n (%)	0	7 (17.1%)	5 (12.8%)	3 (11.1%)	2 (25.0%)	17 (100.0%)	Chi ² =1.242, p=0.743, V=0.104
	1	34 (82.9%)	34 (87.2%)	24 (88.9%)	6 (75.0%)	98 (100.0%)	
Elevated mood, n (%)	0	41 (100.0%)	39 (100.0%)	26 (96.3%)	8 (100.0%)	114 (100.0%)	Chi ² =3.288, p=0.349, V=0.169
	1	0 (0.0%)	0 (0.0%)	1 (3.7%)	0 (0.0%)	1 (100.0%)	
Apathy, n (%)	0	5 (12.2%)	8 (20.5%)	10 (37.0%)	0 (0.0%)	23 (100.0%)	Chi ² =8.466, p=0.037, V=0.271
	1	36 (87.8%)	31 (79.5%)	17 (63.0%)	8 (100.0%)	92 (100.0%)	
Irritability, n (%)	0	32 (78.0%)	34 (87.2%)	19 (70.4%)	8 (100.0%)	93 (100.0%)	Chi ² =5.031, p=0.170, V=0.209
	1	9 (22.0%)	5 (12.8%)	8 (29.6%)	0 (0.0%)	22 (100.0%)	
Anhedonia, n (%)	0	34 (82.9%)	16 (41.0%)	7 (25.9%)	8 (100.0%)	65 (100.0%)	Chi ² =31.882, p=0.000, V=0.527
	1	7 (17.1%)	23 (59.0%)	20 (74.1%)	0 (0.0%)	50 (100.0%)	
Euphoria, n (%)	0	41 (100.0%)	39 (100.0%)	27 (100.0%)	8 (100.0%)	115 (100.0%)	Chi ² =0.000, p=1.000, V=0.000
Fears (except for ED-related symptoms), n (%)	0	39 (95.1%)	39 (100.0%)	27 (100.0%)	8 (100.0%)	113 (100.0%)	Chi ² =3.674, p=0.299, V=0.179
	1	2 (4.9%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	2 (100.0%)	
Somatoform autonomic dysfunction, n (%)	0	40 (97.6%)	39 (100.0%)	23 (85.2%)	1 (12.5%)	103 (100.0%)	Chi ² =58.740, p=0.000, V=0.715
	1	1 (2.4%)	0 (0.0%)	4 (14.8%)	7 (87.5%)	12 (100.0%)	
Hypochondriacal concerns, n (%)	0	41 (100.0%)	36 (92.3%)	18 (66.7%)	0 (0.0%)	95 (100.0%)	Chi ² =53.962, p=0.000, V=0.685

	1	0 (0.0%)	3 (7.7%)	9 (33.3%)	8 (100.0%)	20 (100.0%)	
Body dysmorphic disorder (except for concerns regarding thickness of the body or its parts), <i>n</i> (%)	0	41 (100.0%)	39 (100.0%)	26 (96.3%)	8 (100.0%)	114 (100.0%)	Chi ² =3.288, <i>p</i> =0.349, V=0.169
	1	0 (0.0%)	0 (0.0%)	1 (3.7%)	0 (0.0%)	1 (100.0%)	
Pain, <i>n</i> (%)	0	41 (100.0%)	38 (97.4%)	27 (100.0%)	5 (62.5%)	111 (100.0%)	Chi ² =30.129, <i>p</i> =0.000, V=0.512
	1	0 (0.0%)	1 (2.6%)	0 (0.0%)	3 (37.5%)	4 (100.0%)	
Delusional level of anxiety (including regarding EDs), <i>n</i> (%)	0	21 (51.2%)	36 (92.3%)	19 (70.4%)	5 (62.5%)	81 (100.0%)	Chi ² =16.472, <i>p</i> =0.001, V=0.378
	1	20 (48.8%)	3 (7.7%)	8 (29.6%)	3 (37.5%)	34 (100.0%)	
Formal thought disorders (associative process disorders), <i>n</i> (%)	0	35 (85.4%)	25 (64.1%)	13 (48.1%)	6 (75.0%)	79 (100.0%)	Chi ² =11.130, <i>p</i> =0.011, V=0.311
	1	6 (14.6%)	14 (35.9%)	14 (51.9%)	2 (25.0%)	36 (100.0%)	
Delusional level of depression, <i>n</i> (%)	0	41 (100.0%)	39 (100.0%)	26 (96.3%)	8 (100.0%)	114 (100.0%)	Chi ² =3.288, <i>p</i> =0.349, V=0.169
	1	0 (0.0%)	0 (0.0%)	1 (3.7%)	0 (0.0%)	1 (100.0%)	
Delusional ideas (except for EDs), <i>n</i> (%)	0	38 (92.7%)	39 (100.0%)	25 (92.6%)	6 (75.0%)	108 (100.0%)	Chi ² =7.725, <i>p</i> =0.052, V=0.259
	1	3 (7.3%)	0 (0.0%)	2 (7.4%)	2 (25.0%)	7 (100.0%)	
Overvalued ideas (except for EDs), <i>n</i> (%)	0	41 (100.0%)	39 (100.0%)	27 (100.0%)	7 (87.5%)	114 (100.0%)	Chi ² =13.492, <i>p</i> =0.004, V=0.343
	1	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (12.5%)	1 (100.0%)	
Dysmorphia (including regarding thickness of the body or its parts), <i>n</i> (%)	0	38 (92.7%)	39 (100.0%)	25 (92.6%)	7 (87.5%)	109 (100.0%)	Chi ² =3.632, <i>p</i> =0.304, V=0.178
	1	3 (7.3%)	0 (0.0%)	2 (7.4%)	1 (12.5%)	6 (100.0%)	
Ideas of reference (except for EDs), <i>n</i> (%)	0	40 (97.6%)	39 (100.0%)	27 (100.0%)	6 (75.0%)	112 (100.0%)	Chi ² =17.560, <i>p</i> =0.001, V=0.391
	1	1 (2.4%)	0 (0.0%)	0 (0.0%)	2 (25.0%)	3 (100.0%)	
Reduced attentional focus, <i>n</i> (%)	0	38 (92.7%)	33 (84.6%)	17 (63.0%)	8 (100.0%)	96 (100.0%)	Chi ² =12.378, <i>p</i> =0.006, V=0.328
	1	3 (7.3%)	6 (15.4%)	10 (37.0%)	0 (0.0%)	19 (100.0%)	
Delirium, <i>n</i> (%)	0	41 (100.0%)	39 (100.0%)	27 (100.0%)	8 (100.0%)	115 (100.0%)	Chi ² =0.000, <i>p</i> =1.000, V=0.000
Memory deficits, <i>n</i> (%)	0	41 (100.0%)	39 (100.0%)	22 (81.5%)	8 (100.0%)	110 (100.0%)	Chi ² =17.037, <i>p</i> =0.001, V=0.385
	1	0 (0.0%)	0 (0.0%)	5 (18.5%)	0 (0.0%)	5 (100.0%)	
Impulsive actions, <i>n</i> (%)	0	40 (97.6%)	39 (100.0%)	25 (92.6%)	8 (100.0%)	112 (100.0%)	Chi ² =3.711, <i>p</i> =0.294, V=0.180
	1	1 (2.4%)	0 (0.0%)	2 (7.4%)	0 (0.0%)	3 (100.0%)	
Conflict behavior, <i>n</i> (%)	0	40 (97.6%)	38 (97.4%)	25 (92.6%)	8 (100.0%)	111 (100.0%)	Chi ² =1.759, <i>p</i> =0.624, V=0.124
	1	1 (2.4%)	1 (2.6%)	2 (7.4%)	0 (0.0%)	4 (100.0%)	
Self-harm, <i>n</i> (%)	0	40 (97.6%)	38 (97.4%)	27 (100.0%)	8 (100.0%)	113 (100.0%)	Chi ² =0.892, <i>p</i> =0.827, V=0.088
	1	1 (2.4%)	1 (2.6%)	0 (0.0%)	0 (0.0%)	2 (100.0%)	

Suicidal thoughts, <i>n</i> (%)	0	41 (100.0%)	36 (92.3%)	26 (96.3%)	8 (100.0%)	111 (100.0%)	Chi ² =3.833, <i>p</i> =0.280, V=0.183
	1	0 (0.0%)	3 (7.7%)	1 (3.7%)	0 (0.0%)	4 (100.0%)	
Cenesthopathies, <i>n</i> (%)	0	40 (97.6%)	39 (100.0%)	27 (100.0%)	6 (75.0%)	112 (100.0%)	Chi ² =17.560, <i>p</i> =0.001, V=0.391
	1	1 (2.4%)	0 (0.0%)	0 (0.0%)	2 (25.0%)	3 (100.0%)	
Hallucinations, <i>n</i> (%)	0	39 (95.1%)	39 (100.0%)	26 (96.3%)	7 (87.5%)	111 (100.0%)	Chi ² =3.588, <i>p</i> =0.309, V=0.177
	1	2 (4.9%)	0 (0.0%)	1 (3.7%)	1 (12.5%)	4 (100.0%)	
Depersonalization, <i>n</i> (%)	0	41 (100.0%)	39 (100.0%)	27 (100.0%)	8 (100.0%)	115 (100.0%)	Chi ² =0.000, <i>p</i> =1.000, V=0.000
Obsessive thoughts (except for EDs), <i>n</i> (%)	0	41 (100.0%)	36 (92.3%)	24 (88.9%)	6 (75.0%)	107 (100.0%)	Chi ² =7.842, <i>p</i> =0.049, V=0.261
	1	0 (0.0%)	3 (7.7%)	3 (11.1%)	2 (25.0%)	8 (100.0%)	
Rituals (except for EDs), <i>n</i> (%)	0	41 (100.0%)	38 (97.4%)	25 (92.6%)	8 (100.0%)	112 (100.0%)	Chi ² =3.760, <i>p</i> =0.289, V=0.181
	1	0 (0.0%)	1 (2.6%)	2 (7.4%)	0 (0.0%)	3 (100.0%)	
Compulsions (except for ED-related symptoms), <i>n</i> (%)	0	41 (100.0%)	38 (97.4%)	27 (100.0%)	8 (100.0%)	114 (100.0%)	Chi ² =1.966, <i>p</i> =0.580, V=0.131
	1	0 (0.0%)	1 (2.6%)	0 (0.0%)	0 (0.0%)	1 (100.0%)	
Difficulties falling asleep, <i>n</i> (%)	0	30 (73.2%)	4 (10.3%)	0 (0.0%)	3 (37.5%)	37 (100.0%)	Chi ² =53.075, <i>p</i> =0.000, V=0.679
	1	11 (26.8%)	35 (89.7%)	27 (100.0%)	5 (62.5%)	78 (100.0%)	
Fatigue, <i>n</i> (%)	0	40 (97.6%)	39 (100.0%)	27 (100.0%)	8 (100.0%)	114 (100.0%)	Chi ² =1.821, <i>p</i> =0.610, V=0.126
	1	1 (2.4%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (100.0%)	
Superficial sleep, <i>n</i> (%)	0	35 (85.4%)	23 (59.0%)	14 (51.9%)	8 (100.0%)	80 (100.0%)	Chi ² =14.402, <i>p</i> =0.002, V=0.354
	1	6 (14.6%)	16 (41.0%)	13 (48.1%)	0 (0.0%)	35 (100.0%)	
Early awakening, <i>n</i> (%)	0	40 (97.6%)	39 (100.0%)	26 (96.3%)	8 (100.0%)	113 (100.0%)	Chi ² =1.559, <i>p</i> =0.669, V=0.116
	1	1 (2.4%)	0 (0.0%)	1 (3.7%)	0 (0.0%)	2 (100.0%)	
Decreased appetite, <i>n</i> (%)	0	41 (100.0%)	38 (97.4%)	27 (100.0%)	8 (100.0%)	114 (100.0%)	Chi ² =1.966, <i>p</i> =0.580, V=0.131
	1	0 (0.0%)	1 (2.6%)	0 (0.0%)	0 (0.0%)	1 (100.0%)	
Fears related to eating, weight gain, loss of control, <i>n</i> (%)	0	10 (24.4%)	38 (97.4%)	4 (14.8%)	7 (87.5%)	59 (100.0%)	Chi ² =63.694, <i>p</i> =0.000, V=0.744
	1	31 (75.6%)	1 (2.6%)	23 (85.2%)	1 (12.5%)	56 (100.0%)	
Obsessive ideas (regarding food, weight, and body shape), <i>n</i> (%)	0	33 (80.5%)	37 (94.9%)	16 (59.3%)	8 (100.0%)	94 (100.0%)	Chi ² =15.478, <i>p</i> =0.001, V=0.367
	1	8 (19.5%)	2 (5.1%)	11 (40.7%)	0 (0.0%)	21 (100.0%)	
Body dysmorphic disorder (concerns regarding thickness of the body or its parts), <i>n</i> (%)	0	22 (53.7%)	38 (97.4%)	8 (29.6%)	8 (100.0%)	76 (100.0%)	Chi ² =40.044, <i>p</i> =0.000, V=0.590

	1	19 (46.3%)	1 (2.6%)	19 (70.4%)	0 (0.0%)	39 (100.0%)	
Rituals (food, evaluation of one's own body), <i>n</i> (%)	0	40 (97.6%)	38 (97.4%)	27 (100.0%)	8 (100.0%)	113 (100.0%)	Chi ² =0.892, <i>p</i> =0.827, V=0.088
	1	1 (2.4%)	1 (2.6%)	0 (0.0%)	0 (0.0%)	2 (100.0%)	
Vomiting, <i>n</i> (%)	0	41 (100.0%)	39 (100.0%)	22 (81.5%)	8 (100.0%)	110 (100.0%)	Chi ² =17.037, <i>p</i> =0.001, V=0.385
	1	0 (0.0%)	0 (0.0%)	5 (18.5%)	0 (0.0%)	5 (100.0%)	
Sensitive ideas of reference ("others are critically judging me"), <i>n</i> (%)	0	31 (75.6%)	39 (100.0%)	25 (92.6%)	8 (100.0%)	103 (100.0%)	Chi ² =14.284, <i>p</i> =0.003, V=0.352
	1	10 (24.4%)	0 (0.0%)	2 (7.4%)	0 (0.0%)	12 (100.0%)	
Overvalued ideas (regarding diet, weight, and body shape), <i>n</i> (%)	0	35 (85.4%)	39 (100.0%)	13 (48.1%)	8 (100.0%)	95 (100.0%)	Chi ² =32.429, <i>p</i> =0.000, V=0.531
	1	6 (14.6%)	0 (0.0%)	14 (51.9%)	0 (0.0%)	20 (100.0%)	
Compulsions as part of EDs, <i>n</i> (%)	0	40 (97.6%)	39 (100.0%)	26 (96.3%)	8 (100.0%)	113 (100.0%)	Chi ² =1.559, <i>p</i> =0.669, V=0.116
	1	1 (2.4%)	0 (0.0%)	1 (3.7%)	0 (0.0%)	2 (100.0%)	
Extrapyramidal effects, <i>n</i> (%)	0	40 (97.6%)	35 (89.7%)	15 (55.6%)	6 (75.0%)	96 (100.0%)	Chi ² =22.686, <i>p</i> =0.000, V=0.444
	1	1 (2.4%)	4 (10.3%)	12 (44.4%)	2 (25.0%)	19 (100.0%)	
Akathisia, <i>n</i> (%)	0	41 (100.0%)	36 (92.3%)	26 (96.3%)	7 (87.5%)	110 (100.0%)	Chi ² =4.218, <i>p</i> =0.239, V=0.192
	1	0 (0.0%)	3 (7.7%)	1 (3.7%)	1 (12.5%)	5 (100.0%)	
Prevention of EPS, <i>n</i> (%)	0	26 (63.4%)	36 (92.3%)	15 (55.6%)	5 (62.5%)	82 (100.0%)	Chi ² =13.232, <i>p</i> =0.004, V=0.339
	1	15 (36.6%)	3 (7.7%)	12 (44.4%)	3 (37.5%)	33 (100.0%)	
Results of the M.I.N.I. (<i>n</i>=110)							
Major depressive episode, current (2 weeks), <i>n</i> (%)	0	28 (70.0%)	30 (81.1%)	18 (72.0%)	7 (87.5%)	83 (100.0%)	Chi ² =2.063, <i>p</i> =0.559, V=0.137
	1	12 (30.0%)	7 (18.9%)	7 (28.0%)	1 (12.5%)	27 (100.0%)	
Major depressive episode, recurrent, <i>n</i> (%)	0	25 (62.5%)	20 (54.1%)	10 (40.0%)	7 (87.5%)	62 (100.0%)	Chi ² =6.568, <i>p</i> =0.087, V=0.244
	1	15 (37.5%)	17 (45.9%)	15 (60.0%)	1 (12.5%)	48 (100.0%)	
Major depressive episode, in the past, <i>n</i> (%)	0	35 (87.5%)	34 (91.9%)	23 (92.0%)	7 (87.5%)	99 (100.0%)	Chi ² =0.592, <i>p</i> =0.898, V=0.073
	1	5 (12.5%)	3 (8.1%)	2 (8.0%)	1 (12.5%)	11 (100.0%)	
Current suicidal risk, low, <i>n</i> (%)	0	19 (47.5%)	14 (37.8%)	13 (52.0%)	6 (75.0%)	52 (100.0%)	Chi ² =4.014, <i>p</i> =0.260, V=0.191
	1	21 (52.5%)	23 (62.2%)	12 (48.0%)	2 (25.0%)	58 (100.0%)	
Current suicidal risk, moderate, <i>n</i> (%)	0	37 (92.5%)	35 (94.6%)	19 (76.0%)	8 (100.0%)	99 (100.0%)	Chi ² =7.479, <i>p</i> =0.058, V=0.261
	1	3 (7.5%)	2 (5.4%)	6 (24.0%)	0 (0.0%)	11 (100.0%)	
Current suicidal risk, high, <i>n</i> (%)	0	40 (100.0%)	36 (97.3%)	24 (96.0%)	8 (100.0%)	108 (100.0%)	Chi ² =1.718, <i>p</i> =0.633, V=0.125
	1	0 (0.0%)	1 (2.7%)	1 (4.0%)	0 (0.0%)	2 (100.0%)	

Manic episode, current, <i>n</i> (%)	0	40 (100.0%)	37 (100.0%)	25 (100.0%)	8 (100.0%)	110 (100.0%)	Chi ² =0.000, <i>p</i> =1.000, V=0.000
Manic episode, in the past, <i>n</i> (%)	0	40 (100.0%)	36 (97.3%)	25 (100.0%)	8 (100.0%)	109 (100.0%)	Chi ² =1.991, <i>p</i> =0.574, V=0.135
	1	0 (0.0%)	1 (2.7%)	0 (0.0%)	0 (0.0%)	1 (100.0%)	
Type I bipolar affective disorder, current, <i>n</i> (%)	0	40 (100.0%)	37 (100.0%)	25 (100.0%)	8 (100.0%)	110 (100.0%)	Chi ² =0.000, <i>p</i> =1.000, V=0.000
Type I bipolar affective disorder, in the past, <i>n</i> (%)	0	40 (100.0%)	37 (100.0%)	25 (100.0%)	8 (100.0%)	110 (100.0%)	Chi ² =0.000, <i>p</i> =1.000, V=0.000
Type II bipolar affective disorder, current, <i>n</i> (%)	0	39 (97.5%)	35 (94.6%)	24 (96.0%)	8 (100.0%)	106 (100.0%)	Chi ² =0.789, <i>p</i> =0.852, V=0.085
	1	1 (2.5%)	2 (5.4%)	1 (4.0%)	0 (0.0%)	4 (100.0%)	
Type II bipolar affective disorder, in the past, <i>n</i> (%)	0	39 (97.5%)	36 (97.3%)	25 (100.0%)	8 (100.0%)	108 (100.0%)	Chi ² =0.877, <i>p</i> =0.831, V=0.089
	1	1 (2.5%)	1 (2.7%)	0 (0.0%)	0 (0.0%)	2 (100.0%)	
Bipolar affective disorder, unspecified, current, <i>n</i> (%)	0	40 (100.0%)	37 (100.0%)	25 (100.0%)	8 (100.0%)	110 (100.0%)	Chi ² =0.000, <i>p</i> =1.000, V=0.000
Bipolar affective disorder, unspecified, in the past, <i>n</i> (%)	0	40 (100.0%)	37 (100.0%)	25 (100.0%)	8 (100.0%)	110 (100.0%)	Chi ² =0.000, <i>p</i> =1.000, V=0.000
Panic disorder, current (over the past month), <i>n</i> (%)	0	38 (95.0%)	32 (86.5%)	22 (88.0%)	7 (87.5%)	99 (100.0%)	Chi ² =1.785, <i>p</i> =0.618, V=0.127
	1	2 (5.0%)	5 (13.5%)	3 (12.0%)	1 (12.5%)	11 (100.0%)	
Panic disorder, throughout lifetime, <i>n</i> (%)	0	33 (82.5%)	33 (89.2%)	22 (88.0%)	5 (62.5%)	93 (100.0%)	Chi ² =3.943, <i>p</i> =0.268, V=0.189
	1	7 (17.5%)	4 (10.8%)	3 (12.0%)	3 (37.5%)	17 (100.0%)	
Agoraphobia, current, <i>n</i> (%)	0	36 (90.0%)	33 (89.2%)	24 (96.0%)	8 (100.0%)	101 (100.0%)	Chi ² =1.811, <i>p</i> =0.612, V=0.128
	1	4 (10.0%)	4 (10.8%)	1 (4.0%)	0 (0.0%)	9 (100.0%)	
Social phobia (social anxiety disorder), current (over the past month), <i>n</i> (%)	0	38 (95.0%)	33 (89.2%)	25 (100.0%)	8 (100.0%)	104 (100.0%)	Chi ² =3.978, <i>p</i> =0.264, V=0.190
	1	2 (5.0%)	4 (10.8%)	0 (0.0%)	0 (0.0%)	6 (100.0%)	
Social phobia (social anxiety disorder), generalized, <i>n</i> (%)	0	36 (90.0%)	35 (94.6%)	23 (92.0%)	6 (75.0%)	100 (100.0%)	Chi ² =3.134, <i>p</i> =0.371, V=0.169
	1	4 (10.0%)	2 (5.4%)	2 (8.0%)	2 (25.0%)	10 (100.0%)	
Social phobia (social anxiety disorder), non-generalized, <i>n</i> (%)	0	38 (95.0%)	35 (94.6%)	22 (88.0%)	8 (100.0%)	103 (100.0%)	Chi ² =2.058, <i>p</i> =0.560, V=0.137
	1	2 (5.0%)	2 (5.4%)	3 (12.0%)	0 (0.0%)	7 (100.0%)	
Obsessive-compulsive disorder, current (over the past month), <i>n</i> (%)	0	21 (52.5%)	23 (62.2%)	14 (56.0%)	2 (25.0%)	60 (100.0%)	Chi ² =3.771, <i>p</i> =0.287, V=0.185
	1	19 (47.5%)	14 (37.8%)	11 (44.0%)	6 (75.0%)	50 (100.0%)	
Post-traumatic stress disorder, current (over the past month), <i>n</i> (%)	0	40 (100.0%)	34 (91.9%)	23 (92.0%)	7 (87.5%)	104 (100.0%)	Chi ² =3.897, <i>p</i> =0.273, V=0.188
	1	0 (0.0%)	3 (8.1%)	2 (8.0%)	1 (12.5%)	6 (100.0%)	

Alcohol dependence, over the past 12 months, <i>n</i> (%)	0	38 (95.0%)	33 (89.2%)	23 (92.0%)	8 (100.0%)	102 (100.0%)	Chi ² =1.640, <i>p</i> =0.650, V=0.122
	1	2 (5.0%)	4 (10.8%)	2 (8.0%)	0 (0.0%)	8 (100.0%)	
Alcohol abuse, over the past 12 months, <i>n</i> (%)	0	38 (95.0%)	34 (91.9%)	22 (88.0%)	8 (100.0%)	102 (100.0%)	Chi ² =1.801, <i>p</i> =0.615, V=0.128
	1	2 (5.0%)	3 (8.1%)	3 (12.0%)	0 (0.0%)	8 (100.0%)	
Dependence on psychoactive substances (other than alcohol), over the past 12 months, <i>n</i> (%)	0	40 (100.0%)	35 (94.6%)	24 (96.0%)	8 (100.0%)	107 (100.0%)	Chi ² =2.499, <i>p</i> =0.476, V=0.151
	1	0 (0.0%)	2 (5.4%)	1 (4.0%)	0 (0.0%)	3 (100.0%)	
Psychoactive substances abuse (other than alcohol), over the past 12 months, <i>n</i> (%)	0	40 (100.0%)	35 (94.6%)	24 (96.0%)	8 (100.0%)	107 (100.0%)	Chi ² =2.499, <i>p</i> =0.476, V=0.151
	1	0 (0.0%)	2 (5.4%)	1 (4.0%)	0 (0.0%)	3 (100.0%)	
Psychotic disorders, throughout lifetime, <i>n</i> (%)	0	40 (100.0%)	36 (97.3%)	24 (96.0%)	7 (87.5%)	107 (100.0%)	Chi ² =4.154, <i>p</i> =0.245, V=0.194
	1	0 (0.0%)	1 (2.7%)	1 (4.0%)	1 (12.5%)	3 (100.0%)	
Psychotic disorders, current, <i>n</i> (%)	0	39 (97.5%)	36 (97.3%)	25 (100.0%)	6 (75.0%)	106 (100.0%)	Chi ² =11.603, <i>p</i> =0.009, V=0.325
	1	1 (2.5%)	1 (2.7%)	0 (0.0%)	2 (25.0%)	4 (100.0%)	
Mood disorder with psychotic features, throughout lifetime, <i>n</i> (%)	0	40 (100.0%)	36 (97.3%)	25 (100.0%)	8 (100.0%)	109 (100.0%)	Chi ² =1.991, <i>p</i> =0.574, V=0.135
	1	0 (0.0%)	1 (2.7%)	0 (0.0%)	0 (0.0%)	1 (100.0%)	
Mood disorder with psychotic features, current, <i>n</i> (%)	0	39 (97.5%)	37 (100.0%)	25 (100.0%)	8 (100.0%)	109 (100.0%)	Chi ² =1.766, <i>p</i> =0.622, V=0.127
	1	1 (2.5%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (100.0%)	
Anorexia nervosa, current (over the past 3 months), <i>n</i> (%)	0	8 (20.0%)	9 (24.3%)	8 (32.0%)	1 (12.5%)	26 (100.0%)	Chi ² =1.821, <i>p</i> =0.610, V=0.129
	1	32 (80.0%)	28 (75.7%)	17 (68.0%)	7 (87.5%)	84 (100.0%)	
Bulimia nervosa, current (over the past 3 months), <i>n</i> (%)	0	40 (100.0%)	37 (100.0%)	25 (100.0%)	8 (100.0%)	110 (100.0%)	Chi ² =0.000, <i>p</i> =1.000, V=0.000
Anorexia nervosa, binge-purge pattern, current, <i>n</i> (%)	0	29 (72.5%)	26 (70.3%)	11 (44.0%)	7 (87.5%)	73 (100.0%)	Chi ² =8.130, <i>p</i> =0.043, V=0.272
	1	11 (27.5%)	11 (29.7%)	14 (56.0%)	1 (12.5%)	37 (100.0%)	
Generalized anxiety disorder, current (over the past 6 months), <i>n</i> (%)	0	27 (67.5%)	24 (64.9%)	16 (64.0%)	4 (50.0%)	71 (100.0%)	Chi ² =0.897, <i>p</i> =0.826, V=0.090
	1	13 (32.5%)	13 (35.1%)	9 (36.0%)	4 (50.0%)	39 (100.0%)	
The presence of medical, structural drug-related causes, <i>n</i> (%)	0	39 (97.5%)	35 (94.6%)	22 (88.0%)	8 (100.0%)	104 (100.0%)	Chi ² =3.216, <i>p</i> =0.360, V=0.171
	1	1 (2.5%)	2 (5.4%)	3 (12.0%)	0 (0.0%)	6 (100.0%)	
SCL-90-R							
Somatization (Median [Q1; Q3])		0.71 [0.31; 1.08]	0.75 [0.42; 1.50]	1.17 [0.58; 1.67]	1.08 [0.83; 1.46]	0.83 [0.50; 1.40]	Kruskal: H=5.747, <i>p</i> =0.125, ε ² =0.053
Obsessions (Median [Q1; Q3])		1.20 [0.67; 2.12]	1.40 [0.50; 1.90]	1.90 [1.30; 2.50]	1.45 [1.10; 2.05]	1.40 [0.70; 2.10]	Kruskal: H=4.714, <i>p</i> =0.194, ε ² =0.043
Sensitivity (Median [Q1; Q3])		1.27 [0.64; 2.00]	1.11 [0.56; 2.00]	1.67 [1.00; 2.33]	1.00 [0.86; 1.19]	1.33 [0.59; 2.00]	Kruskal: H=3.422, <i>p</i> =0.331, ε ² =0.031

Depression (Median [Q1; Q3])		1.38 [0.90; 1.87]	1.46 [0.62; 2.08]	1.85 [0.92; 2.54]	1.47 [1.25; 1.71]	1.50 [0.92; 2.13]	Kruskal: $H=2.584$, $p=0.460$, $\epsilon^2=0.024$
Trait anxiety (Median [Q1; Q3])		0.80 [0.35; 1.32]	0.70 [0.30; 1.70]	1.10 [0.50; 2.10]	1.30 [1.00; 1.30]	0.90 [0.40; 1.60]	Kruskal: $H=3.326$, $p=0.344$, $\epsilon^2=0.031$
Hostility (Median [Q1; Q3])		0.75 [0.17; 1.17]	0.50 [0.17; 1.33]	0.67 [0.33; 1.67]	0.33 [0.17; 0.54]	0.67 [0.17; 1.33]	Kruskal: $H=4.737$, $p=0.192$, $\epsilon^2=0.043$
Phobia (Median [Q1; Q3])		0.43 [0.00; 1.00]	0.29 [0.00; 0.57]	0.43 [0.00; 1.00]	0.57 [0.25; 0.86]	0.43 [0.00; 0.86]	Kruskal: $H=2.152$, $p=0.542$, $\epsilon^2=0.020$
Paranoid ideation (Median [Q1; Q3])		0.42 [0.17; 1.17]	0.67 [0.17; 1.17]	1.17 [0.33; 1.50]	0.58 [0.46; 1.04]	0.67 [0.17; 1.17]	Kruskal: $H=4.559$, $p=0.207$, $\epsilon^2=0.042$
Psychoticism (Median [Q1; Q3])		0.55 [0.40; 1.10]	0.50 [0.30; 1.40]	0.80 [0.40; 1.50]	0.55 [0.15; 0.75]	0.60 [0.30; 1.27]	Kruskal: $H=2.142$, $p=0.544$, $\epsilon^2=0.020$
Global Severity Index (Median [Q1; Q3])		0.95 [0.61; 1.50]	1.02 [0.41; 1.37]	1.27 [0.96; 1.94]	1.09 [0.83; 1.15]	1.02 [0.61; 1.58]	Kruskal: $H=3.832$, $p=0.280$, $\epsilon^2=0.035$
Positive Symptom Total (Median [Q1; Q3])		46.50 [31.00; 60.25]	44.00 [22.00; 61.00]	62.00 [45.00; 73.00]	57.50 [36.50; 61.25]	50.00 [31.00; 64.00]	Kruskal: $H=5.744$, $p=0.125$, $\epsilon^2=0.053$
Positive Symptom Distress Index (Median [Q1; Q3])		1.96 [1.49; 2.28]	1.91 [1.50; 2.30]	2.12 [1.48; 2.42]	1.69 [1.59; 1.96]	1.92 [1.49; 2.30]	Kruskal: $H=1.329$, $p=0.722$, $\epsilon^2=0.012$

Note: BMI — Body Mass Index; Kruskal — the Kruskal–Wallis H test; M.I.N.I. — Mini-International Neuropsychiatric Interview; SD — mean value.

ПРИЛОЖЕНИЕ

Таблица П2. Сравнительная характеристика кластеров по всем параметрам исследования

Переменная	Ответ на категориальную переменную	Кластер 1 (n=41)	Кластер 2 (n=39)	Кластер 3 (n=27)	Кластер 4 (n=8)	Всего (n=115)	Тест
Min of ИМТ (Median [Q1; Q3])	—	13,90 [12,60; 15,65]	15,40 [14,10; 16,70]	14,00 [11,65; 16,45]	14,50 [13,62; 15,75]	14,57 [12,65; 16,30]	Kruskal: H=7,429, p=0,059, ε ² =0,065
Возраст (Median [Q1; Q3])	—	21,00 [18,00; 24,00]	22,00 [19,25; 26,00]	27,00 [21,00; 31,00]	21,50 [20,75; 29,50]	22,00 [19,00; 26,00]	Kruskal: H=9,482, p=0,024, ε ² =0,086
ИМТ (Mean (SD))	—	14,16 (2,02)	15,45 (1,91)	14,14 (2,68)	14,71 (2,13)	14,63 (2,21)	ANOVA: F=2,825, p=0,042, d=0,593
Длительность заболевания (лет) (Median [Q1; Q3])	—	4,00 [1,00; 7,50]	5,00 [3,00; 9,00]	6,00 [3,00; 9,00]	3,50 [1,75; 5,50]	5,00 [2,00; 9,00]	Kruskal: H=5,459, p=0,141, ε ² =0,050
Чек-лист симптомов (n=115)							
Тоска, n (%)	0	37 (90,2%)	16 (41,0%)	3 (11,1%)	5 (62,5%)	61 (100,0%)	Chi2=44,389, p=0,000, V=0,621
	1	4 (9,8%)	23 (59,0%)	24 (88,9%)	3 (37,5%)	54 (100,0%)	
Перепады настроения, n (%)	0	26 (63,4%)	26 (66,7%)	12 (44,4%)	7 (87,5%)	71 (100,0%)	Chi2=6,116, p=0,106, V=0,231
	1	15 (36,6%)	13 (33,3%)	15 (55,6%)	1 (12,5%)	44 (100,0%)	
Тревога, n (%)	0	7 (17,1%)	5 (12,8%)	3 (11,1%)	2 (25,0%)	17 (100,0%)	Chi2=1,242, p=0,743, V=0,104
	1	34 (82,9%)	34 (87,2%)	24 (88,9%)	6 (75,0%)	98 (100,0%)	
Приподнятое настроение, n (%)	0	41 (100,0%)	39 (100,0%)	26 (96,3%)	8 (100,0%)	114 (100,0%)	Chi2=3,288, p=0,349, V=0,169
	1	0 (0,0%)	0 (0,0%)	1 (3,7%)	0 (0,0%)	1 (100,0%)	
Апатия, n (%)	0	5 (12,2%)	8 (20,5%)	10 (37,0%)	0 (0,0%)	23 (100,0%)	Chi2=8,466, p=0,037, V=0,271
	1	36 (87,8%)	31 (79,5%)	17 (63,0%)	8 (100,0%)	92 (100,0%)	
Раздражительность, n (%)	0	32 (78,0%)	34 (87,2%)	19 (70,4%)	8 (100,0%)	93 (100,0%)	Chi2=5,031, p=0,170, V=0,209
	1	9 (22,0%)	5 (12,8%)	8 (29,6%)	0 (0,0%)	22 (100,0%)	
Ангедония, n (%)	0	34 (82,9%)	16 (41,0%)	7 (25,9%)	8 (100,0%)	65 (100,0%)	Chi2=31,882, p=0,000, V=0,527
	1	7 (17,1%)	23 (59,0%)	20 (74,1%)	0 (0,0%)	50 (100,0%)	
Эйфория, n (%)	0	41 (100,0%)	39 (100,0%)	27 (100,0%)	8 (100,0%)	115 (100,0%)	Chi2=0,000, p=1,000, V=0,000
Страхи (кроме симптомов РПП), n (%)	0	39 (95,1%)	39 (100,0%)	27 (100,0%)	8 (100,0%)	113 (100,0%)	Chi2=3,674, p=0,299, V=0,179
	1	2 (4,9%)	0 (0,0%)	0 (0,0%)	0 (0,0%)	2 (100,0%)	

Соматоформная вегетативная дисфункция, <i>n</i> (%)	0	40 (97,6%)	39 (100,0%)	23 (85,2%)	1 (12,5%)	103 (100,0%)	Chi2=58,740, <i>p</i> =0,000, <i>V</i> =0,715
	1	1 (2,4%)	0 (0,0%)	4 (14,8%)	7 (87,5%)	12 (100,0%)	
Ипохондрические переживания, <i>n</i> (%)	0	41 (100,0%)	36 (92,3%)	18 (66,7%)	0 (0,0%)	95 (100,0%)	Chi2=53,962, <i>p</i> =0,000, <i>V</i> =0,685
	1	0 (0,0%)	3 (7,7%)	9 (33,3%)	8 (100,0%)	20 (100,0%)	
Дисморфофобия (кроме переживаний в отношении толщины тела или его частей), <i>n</i> (%)	0	41 (100,0%)	39 (100,0%)	26 (96,3%)	8 (100,0%)	114 (100,0%)	Chi2=3,288, <i>p</i> =0,349, <i>V</i> =0,169
	1	0 (0,0%)	0 (0,0%)	1 (3,7%)	0 (0,0%)	1 (100,0%)	
Боли, <i>n</i> (%)	0	41 (100,0%)	38 (97,4%)	27 (100,0%)	5 (62,5%)	111 (100,0%)	Chi2=30,129, <i>p</i> =0,000, <i>V</i> =0,512
	1	0 (0,0%)	1 (2,6%)	0 (0,0%)	3 (37,5%)	4 (100,0%)	
Бредовый уровень тревоги (в том числе в отношении РПП), <i>n</i> (%)	0	21 (51,2%)	36 (92,3%)	19 (70,4%)	5 (62,5%)	81 (100,0%)	Chi2=16,472, <i>p</i> =0,001, <i>V</i> =0,378
	1	20 (48,8%)	3 (7,7%)	8 (29,6%)	3 (37,5%)	34 (100,0%)	
Формальные нарушения мышления (расстройства ассоциативного процесса), <i>n</i> (%)	0	35 (85,4%)	25 (64,1%)	13 (48,1%)	6 (75,0%)	79 (100,0%)	Chi2=11,130, <i>p</i> =0,011, <i>V</i> =0,311
	1	6 (14,6%)	14 (35,9%)	14 (51,9%)	2 (25,0%)	36 (100,0%)	
Бредовый уровень депрессии, <i>n</i> (%)	0	41 (100,0%)	39 (100,0%)	26 (96,3%)	8 (100,0%)	114 (100,0%)	Chi2=3,288, <i>p</i> =0,349, <i>V</i> =0,169
	1	0 (0,0%)	0 (0,0%)	1 (3,7%)	0 (0,0%)	1 (100,0%)	
Бредовые идеи (кроме РПП), <i>n</i> (%)	0	38 (92,7%)	39 (100,0%)	25 (92,6%)	6 (75,0%)	108 (100,0%)	Chi2=7,725, <i>p</i> =0,052, <i>V</i> =0,259
	1	3 (7,3%)	0 (0,0%)	2 (7,4%)	2 (25,0%)	7 (100,0%)	
Сверхценные идеи (кроме РПП), <i>n</i> (%)	0	41 (100,0%)	39 (100,0%)	27 (100,0%)	7 (87,5%)	114 (100,0%)	Chi2=13,492, <i>p</i> =0,004, <i>V</i> =0,343
	1	0 (0,0%)	0 (0,0%)	0 (0,0%)	1 (12,5%)	1 (100,0%)	
Дисморфомания (в том числе и в отношении толщины тела или его частей), <i>n</i> (%)	0	38 (92,7%)	39 (100,0%)	25 (92,6%)	7 (87,5%)	109 (100,0%)	Chi2=3,632, <i>p</i> =0,304, <i>V</i> =0,178
	1	3 (7,3%)	0 (0,0%)	2 (7,4%)	1 (12,5%)	6 (100,0%)	
Идеи отношения (кроме РПП), <i>n</i> (%)	0	40 (97,6%)	39 (100,0%)	27 (100,0%)	6 (75,0%)	112 (100,0%)	Chi2=17,560, <i>p</i> =0,001, <i>V</i> =0,391
	1	1 (2,4%)	0 (0,0%)	0 (0,0%)	2 (25,0%)	3 (100,0%)	
Снижение концентрации внимания, <i>n</i> (%)	0	38 (92,7%)	33 (84,6%)	17 (63,0%)	8 (100,0%)	96 (100,0%)	Chi2=12,378, <i>p</i> =0,006, <i>V</i> =0,328
	1	3 (7,3%)	6 (15,4%)	10 (37,0%)	0 (0,0%)	19 (100,0%)	
Делирий, <i>n</i> (%)	0	41 (100,0%)	39 (100,0%)	27 (100,0%)	8 (100,0%)	115 (100,0%)	Chi2=0,000, <i>p</i> =1,000, <i>V</i> =0,000
Нарушение памяти, <i>n</i> (%)	0	41 (100,0%)	39 (100,0%)	22 (81,5%)	8 (100,0%)	110 (100,0%)	Chi2=17,037, <i>p</i> =0,001, <i>V</i> =0,385
	1	0 (0,0%)	0 (0,0%)	5 (18,5%)	0 (0,0%)	5 (100,0%)	
Импульсивные действия, <i>n</i> (%)	0	40 (97,6%)	39 (100,0%)	25 (92,6%)	8 (100,0%)	112 (100,0%)	Chi2=3,711, <i>p</i> =0,294, <i>V</i> =0,180
	1	1 (2,4%)	0 (0,0%)	2 (7,4%)	0 (0,0%)	3 (100,0%)	

Конфликтное поведение, <i>n</i> (%)	0	40 (97,6%)	38 (97,4%)	25 (92,6%)	8 (100,0%)	111 (100,0%)	Chi2=1,759, <i>p</i> =0,624, <i>V</i> =0,124
	1	1 (2,4%)	1 (2,6%)	2 (7,4%)	0 (0,0%)	4 (100,0%)	
Самоповреждения, <i>n</i> (%)	0	40 (97,6%)	38 (97,4%)	27 (100,0%)	8 (100,0%)	113 (100,0%)	Chi2=0,892, <i>p</i> =0,827, <i>V</i> =0,088
	1	1 (2,4%)	1 (2,6%)	0 (0,0%)	0 (0,0%)	2 (100,0%)	
Суицидальные мысли, <i>n</i> (%)	0	41 (100,0%)	36 (92,3%)	26 (96,3%)	8 (100,0%)	111 (100,0%)	Chi2=3,833, <i>p</i> =0,280, <i>V</i> =0,183
	1	0 (0,0%)	3 (7,7%)	1 (3,7%)	0 (0,0%)	4 (100,0%)	
Сенестопатии, <i>n</i> (%)	0	40 (97,6%)	39 (100,0%)	27 (100,0%)	6 (75,0%)	112 (100,0%)	Chi2=17,560, <i>p</i> =0,001, <i>V</i> =0,391
	1	1 (2,4%)	0 (0,0%)	0 (0,0%)	2 (25,0%)	3 (100,0%)	
Галлюцинации, <i>n</i> (%)	0	39 (95,1%)	39 (100,0%)	26 (96,3%)	7 (87,5%)	111 (100,0%)	Chi2=3,588, <i>p</i> =0,309, <i>V</i> =0,177
	1	2 (4,9%)	0 (0,0%)	1 (3,7%)	1 (12,5%)	4 (100,0%)	
Деперсонализация, <i>n</i> (%)	0	41 (100,0%)	39 (100,0%)	27 (100,0%)	8 (100,0%)	115 (100,0%)	Chi2=0,000, <i>p</i> =1,000, <i>V</i> =0,000
Навязчивые мысли (кроме РПП), <i>n</i> (%)	0	41 (100,0%)	36 (92,3%)	24 (88,9%)	6 (75,0%)	107 (100,0%)	Chi2=7,842, <i>p</i> =0,049, <i>V</i> =0,261
	1	0 (0,0%)	3 (7,7%)	3 (11,1%)	2 (25,0%)	8 (100,0%)	
Ритуалы (кроме РПП), <i>n</i> (%)	0	41 (100,0%)	38 (97,4%)	25 (92,6%)	8 (100,0%)	112 (100,0%)	Chi2=3,760, <i>p</i> =0,289, <i>V</i> =0,181
	1	0 (0,0%)	1 (2,6%)	2 (7,4%)	0 (0,0%)	3 (100,0%)	
Компульсии (кроме симптомов РПП), <i>n</i> (%)	0	41 (100,0%)	38 (97,4%)	27 (100,0%)	8 (100,0%)	114 (100,0%)	Chi2=1,966, <i>p</i> =0,580, <i>V</i> =0,131
	1	0 (0,0%)	1 (2,6%)	0 (0,0%)	0 (0,0%)	1 (100,0%)	
Нарушение засыпания, <i>n</i> (%)	0	30 (73,2%)	4 (10,3%)	0 (0,0%)	3 (37,5%)	37 (100,0%)	Chi2=53,075, <i>p</i> =0,000, <i>V</i> =0,679
	1	11 (26,8%)	35 (89,7%)	27 (100,0%)	5 (62,5%)	78 (100,0%)	
Утомляемость, <i>n</i> (%)	0	40 (97,6%)	39 (100,0%)	27 (100,0%)	8 (100,0%)	114 (100,0%)	Chi2=1,821, <i>p</i> =0,610, <i>V</i> =0,126
	1	1 (2,4%)	0 (0,0%)	0 (0,0%)	0 (0,0%)	1 (100,0%)	
Поверхностный сон, <i>n</i> (%)	0	35 (85,4%)	23 (59,0%)	14 (51,9%)	8 (100,0%)	80 (100,0%)	Chi2=14,402, <i>p</i> =0,002, <i>V</i> =0,354
	1	6 (14,6%)	16 (41,0%)	13 (48,1%)	0 (0,0%)	35 (100,0%)	
Ранние пробуждения, <i>n</i> (%)	0	40 (97,6%)	39 (100,0%)	26 (96,3%)	8 (100,0%)	113 (100,0%)	Chi2=1,559, <i>p</i> =0,669, <i>V</i> =0,116
	1	1 (2,4%)	0 (0,0%)	1 (3,7%)	0 (0,0%)	2 (100,0%)	
Пониженный аппетит, <i>n</i> (%)	0	41 (100,0%)	38 (97,4%)	27 (100,0%)	8 (100,0%)	114 (100,0%)	Chi2=1,966, <i>p</i> =0,580, <i>V</i> =0,131
	1	0 (0,0%)	1 (2,6%)	0 (0,0%)	0 (0,0%)	1 (100,0%)	
Страхи (еды, набора веса, потери контроля), <i>n</i> (%)	0	10 (24,4%)	38 (97,4%)	4 (14,8%)	7 (87,5%)	59 (100,0%)	Chi2=63,694, <i>p</i> =0,000, <i>V</i> =0,744
	1	31 (75,6%)	1 (2,6%)	23 (85,2%)	1 (12,5%)	56 (100,0%)	
Навязчивые идеи (в отношении питания, веса, фигуры), <i>n</i> (%)	0	33 (80,5%)	37 (94,9%)	16 (59,3%)	8 (100,0%)	94 (100,0%)	Chi2=15,478, <i>p</i> =0,001, <i>V</i> =0,367
	1	8 (19,5%)	2 (5,1%)	11 (40,7%)	0 (0,0%)	21 (100,0%)	
Дисморфофобия (переживания в отношении толщины тела или его частей), <i>n</i> (%)	0	22 (53,7%)	38 (97,4%)	8 (29,6%)	8 (100,0%)	76 (100,0%)	Chi2=40,044, <i>p</i> =0,000, <i>V</i> =0,590
	1	19 (46,3%)	1 (2,6%)	19 (70,4%)	0 (0,0%)	39 (100,0%)	
Ритуалы (еда, оценка своего тела), <i>n</i> (%)	0	40 (97,6%)	38 (97,4%)	27 (100,0%)	8 (100,0%)	113 (100,0%)	Chi2=0,892, <i>p</i> =0,827, <i>V</i> =0,088
	1	1 (2,4%)	1 (2,6%)	0 (0,0%)	0 (0,0%)	2 (100,0%)	

Рвота, <i>n</i> (%)	0	41 (100,0%)	39 (100,0%)	22 (81,5%)	8 (100,0%)	110 (100,0%)	Chi2=17,037, <i>p</i> =0,001, <i>V</i> =0,385
	1	0 (0,0%)	0 (0,0%)	5 (18,5%)	0 (0,0%)	5 (100,0%)	
Сенситивные идеи отношения (другие критически оценивают), <i>n</i> (%)	0	31 (75,6%)	39 (100,0%)	25 (92,6%)	8 (100,0%)	103 (100,0%)	Chi2=14,284, <i>p</i> =0,003, <i>V</i> =0,352
	1	10 (24,4%)	0 (0,0%)	2 (7,4%)	0 (0,0%)	12 (100,0%)	
Сверхценные идеи (в отношении питания, веса, фигуры), <i>n</i> (%)	0	35 (85,4%)	39 (100,0%)	13 (48,1%)	8 (100,0%)	95 (100,0%)	Chi2=32,429, <i>p</i> =0,000, <i>V</i> =0,531
	1	6 (14,6%)	0 (0,0%)	14 (51,9%)	0 (0,0%)	20 (100,0%)	
Компульсии в рамках РПП, <i>n</i> (%)	0	40 (97,6%)	39 (100,0%)	26 (96,3%)	8 (100,0%)	113 (100,0%)	Chi2=1,559, <i>p</i> =0,669, <i>V</i> =0,116
	1	1 (2,4%)	0 (0,0%)	1 (3,7%)	0 (0,0%)	2 (100,0%)	
Экстрапирамидные эффекты, <i>n</i> (%)	0	40 (97,6%)	35 (89,7%)	15 (55,6%)	6 (75,0%)	96 (100,0%)	Chi2=22,686, <i>p</i> =0,000, <i>V</i> =0,444
	1	1 (2,4%)	4 (10,3%)	12 (44,4%)	2 (25,0%)	19 (100,0%)	
Акатизия, <i>n</i> (%)	0	41 (100,0%)	36 (92,3%)	26 (96,3%)	7 (87,5%)	110 (100,0%)	Chi2=4,218, <i>p</i> =0,239, <i>V</i> =0,192
	1	0 (0,0%)	3 (7,7%)	1 (3,7%)	1 (12,5%)	5 (100,0%)	
Профилактика ЭПС, <i>n</i> (%)	0	26 (63,4%)	36 (92,3%)	15 (55,6%)	5 (62,5%)	82 (100,0%)	Chi2=13,232, <i>p</i> =0,004, <i>V</i> =0,339
	1	15 (36,6%)	3 (7,7%)	12 (44,4%)	3 (37,5%)	33 (100,0%)	
Результаты M.I.N.I. (<i>n</i>=110)							
Большой депрессивный эпизод, текущий (2 нед.), <i>n</i> (%)	0	28 (70,0%)	30 (81,1%)	18 (72,0%)	7 (87,5%)	83 (100,0%)	Chi2=2,063, <i>p</i> =0,559, <i>V</i> =0,137
	1	12 (30,0%)	7 (18,9%)	7 (28,0%)	1 (12,5%)	27 (100,0%)	
Большой депрессивный эпизод, рекуррентный, <i>n</i> (%)	0	25 (62,5%)	20 (54,1%)	10 (40,0%)	7 (87,5%)	62 (100,0%)	Chi2=6,568, <i>p</i> =0,087, <i>V</i> =0,244
	1	15 (37,5%)	17 (45,9%)	15 (60,0%)	1 (12,5%)	48 (100,0%)	
Большой депрессивный эпизод, в прошлом, <i>n</i> (%)	0	35 (87,5%)	34 (91,9%)	23 (92,0%)	7 (87,5%)	99 (100,0%)	Chi2=0,592, <i>p</i> =0,898, <i>V</i> =0,073
	1	5 (12,5%)	3 (8,1%)	2 (8,0%)	1 (12,5%)	11 (100,0%)	
Суицидальный риск, текущий, низкий, <i>n</i> (%)	0	19 (47,5%)	14 (37,8%)	13 (52,0%)	6 (75,0%)	52 (100,0%)	Chi2=4,014, <i>p</i> =0,260, <i>V</i> =0,191
	1	21 (52,5%)	23 (62,2%)	12 (48,0%)	2 (25,0%)	58 (100,0%)	
Суицидальный риск, текущий, умеренный, <i>n</i> (%)	0	37 (92,5%)	35 (94,6%)	19 (76,0%)	8 (100,0%)	99 (100,0%)	Chi2=7,479, <i>p</i> =0,058, <i>V</i> =0,261
	1	3 (7,5%)	2 (5,4%)	6 (24,0%)	0 (0,0%)	11 (100,0%)	
Суицидальный риск, текущий, высокий, <i>n</i> (%)	0	40 (100,0%)	36 (97,3%)	24 (96,0%)	8 (100,0%)	108 (100,0%)	Chi2=1,718, <i>p</i> =0,633, <i>V</i> =0,125
	1	0 (0,0%)	1 (2,7%)	1 (4,0%)	0 (0,0%)	2 (100,0%)	
Маниакальный эпизод, текущий, <i>n</i> (%)	0	40 (100,0%)	37 (100,0%)	25 (100,0%)	8 (100,0%)	110 (100,0%)	Chi2=0,000, <i>p</i> =1,000, <i>V</i> =0,000
Маниакальный эпизод, в прошлом, <i>n</i> (%)	0	40 (100,0%)	36 (97,3%)	25 (100,0%)	8 (100,0%)	109 (100,0%)	Chi2=1,991, <i>p</i> =0,574, <i>V</i> =0,135
	1	0 (0,0%)	1 (2,7%)	0 (0,0%)	0 (0,0%)	1 (100,0%)	
Биполярное расстройство I типа, текущее, <i>n</i> (%)	0	40 (100,0%)	37 (100,0%)	25 (100,0%)	8 (100,0%)	110 (100,0%)	Chi2=0,000, <i>p</i> =1,000, <i>V</i> =0,000

Биполярное расстройство I типа, в прошлом, <i>n</i> (%)	0	40 (100,0%)	37 (100,0%)	25 (100,0%)	8 (100,0%)	110 (100,0%)	Chi2=0,000, <i>p</i> =1,000, <i>V</i> =0,000
Биполярное расстройство II типа, текущее, <i>n</i> (%)	0	39 (97,5%)	35 (94,6%)	24 (96,0%)	8 (100,0%)	106 (100,0%)	Chi2=0,789, <i>p</i> =0,852, <i>V</i> =0,085
	1	1 (2,5%)	2 (5,4%)	1 (4,0%)	0 (0,0%)	4 (100,0%)	
Биполярное расстройство II типа, в прошлом, <i>n</i> (%)	0	39 (97,5%)	36 (97,3%)	25 (100,0%)	8 (100,0%)	108 (100,0%)	Chi2=0,877, <i>p</i> =0,831, <i>V</i> =0,089
	1	1 (2,5%)	1 (2,7%)	0 (0,0%)	0 (0,0%)	2 (100,0%)	
Биполярное расстройство неуточненное, текущее, <i>n</i> (%)	0	40 (100,0%)	37 (100,0%)	25 (100,0%)	8 (100,0%)	110 (100,0%)	Chi2=0,000, <i>p</i> =1,000, <i>V</i> =0,000
Биполярное расстройство неуточненное, в прошлом, <i>n</i> (%)	0	40 (100,0%)	37 (100,0%)	25 (100,0%)	8 (100,0%)	110 (100,0%)	Chi2=0,000, <i>p</i> =1,000, <i>V</i> =0,000
Паническое расстройство, текущее (за последний месяц), <i>n</i> (%)	0	38 (95,0%)	32 (86,5%)	22 (88,0%)	7 (87,5%)	99 (100,0%)	Chi2=1,785, <i>p</i> =0,618, <i>V</i> =0,127
	1	2 (5,0%)	5 (13,5%)	3 (12,0%)	1 (12,5%)	11 (100,0%)	
Паническое расстройство, в течение жизни, <i>n</i> (%)	0	33 (82,5%)	33 (89,2%)	22 (88,0%)	5 (62,5%)	93 (100,0%)	Chi2=3,943, <i>p</i> =0,268, <i>V</i> =0,189
	1	7 (17,5%)	4 (10,8%)	3 (12,0%)	3 (37,5%)	17 (100,0%)	
Агорафобия, текущая, <i>n</i> (%)	0	36 (90,0%)	33 (89,2%)	24 (96,0%)	8 (100,0%)	101 (100,0%)	Chi2=1,811, <i>p</i> =0,612, <i>V</i> =0,128
	1	4 (10,0%)	4 (10,8%)	1 (4,0%)	0 (0,0%)	9 (100,0%)	
Социальная фобия (социальное тревожное расстройство), текущая (за последний месяц), <i>n</i> (%)	0	38 (95,0%)	33 (89,2%)	25 (100,0%)	8 (100,0%)	104 (100,0%)	Chi2=3,978, <i>p</i> =0,264, <i>V</i> =0,190
	1	2 (5,0%)	4 (10,8%)	0 (0,0%)	0 (0,0%)	6 (100,0%)	
Социальная фобия (социальное тревожное расстройство), генерализованная, <i>n</i> (%)	0	36 (90,0%)	35 (94,6%)	23 (92,0%)	6 (75,0%)	100 (100,0%)	Chi2=3,134, <i>p</i> =0,371, <i>V</i> =0,169
	1	4 (10,0%)	2 (5,4%)	2 (8,0%)	2 (25,0%)	10 (100,0%)	
Социальная фобия (социальное тревожное расстройство), негенерализованная, <i>n</i> (%)	0	38 (95,0%)	35 (94,6%)	22 (88,0%)	8 (100,0%)	103 (100,0%)	Chi2=2,058, <i>p</i> =0,560, <i>V</i> =0,137
	1	2 (5,0%)	2 (5,4%)	3 (12,0%)	0 (0,0%)	7 (100,0%)	
Обсессивно-компульсивное расстройство, текущее (за последний месяц), <i>n</i> (%)	0	21 (52,5%)	23 (62,2%)	14 (56,0%)	2 (25,0%)	60 (100,0%)	Chi2=3,771, <i>p</i> =0,287, <i>V</i> =0,185
	1	19 (47,5%)	14 (37,8%)	11 (44,0%)	6 (75,0%)	50 (100,0%)	
Посттравматическое стрессовое расстройство, текущее (за последний месяц), <i>n</i> (%)	0	40 (100,0%)	34 (91,9%)	23 (92,0%)	7 (87,5%)	104 (100,0%)	Chi2=3,897, <i>p</i> =0,273, <i>V</i> =0,188
	1	0 (0,0%)	3 (8,1%)	2 (8,0%)	1 (12,5%)	6 (100,0%)	

Зависимость от алкоголя, за последние 12 месяцев, <i>n</i> (%)	0	38 (95,0%)	33 (89,2%)	23 (92,0%)	8 (100,0%)	102 (100,0%)	Chi2=1,640, <i>p</i> =0,650, <i>V</i> =0,122
	1	2 (5,0%)	4 (10,8%)	2 (8,0%)	0 (0,0%)	8 (100,0%)	
Злоупотребление алкоголем, за последние 12 месяцев, <i>n</i> (%)	0	38 (95,0%)	34 (91,9%)	22 (88,0%)	8 (100,0%)	102 (100,0%)	Chi2=1,801, <i>p</i> =0,615, <i>V</i> =0,128
	1	2 (5,0%)	3 (8,1%)	3 (12,0%)	0 (0,0%)	8 (100,0%)	
Зависимость от психоактивных веществ (не алкоголь), за последние 12 месяцев, <i>n</i> (%)	0	40 (100,0%)	35 (94,6%)	24 (96,0%)	8 (100,0%)	107 (100,0%)	Chi2=2,499, <i>p</i> =0,476, <i>V</i> =0,151
	1	0 (0,0%)	2 (5,4%)	1 (4,0%)	0 (0,0%)	3 (100,0%)	
Злоупотребление психоактивными веществами (не алкоголь), за последние 12 месяцев, <i>n</i> (%)	0	40 (100,0%)	35 (94,6%)	24 (96,0%)	8 (100,0%)	107 (100,0%)	Chi2=2,499, <i>p</i> =0,476, <i>V</i> =0,151
	1	0 (0,0%)	2 (5,4%)	1 (4,0%)	0 (0,0%)	3 (100,0%)	
Психотические расстройства, на протяжении всей жизни, <i>n</i> (%)	0	40 (100,0%)	36 (97,3%)	24 (96,0%)	7 (87,5%)	107 (100,0%)	Chi2=4,154, <i>p</i> =0,245, <i>V</i> =0,194
	1	0 (0,0%)	1 (2,7%)	1 (4,0%)	1 (12,5%)	3 (100,0%)	
Психотические расстройства, текущие, <i>n</i> (%)	0	39 (97,5%)	36 (97,3%)	25 (100,0%)	6 (75,0%)	106 (100,0%)	Chi2=11,603, <i>p</i> =0,009, <i>V</i> =0,325
	1	1 (2,5%)	1 (2,7%)	0 (0,0%)	2 (25,0%)	4 (100,0%)	
Расстройство настроения с психотическими симптомами, на протяжении всей жизни, <i>n</i> (%)	0	40 (100,0%)	36 (97,3%)	25 (100,0%)	8 (100,0%)	109 (100,0%)	Chi2=1,991, <i>p</i> =0,574, <i>V</i> =0,135
	1	0 (0,0%)	1 (2,7%)	0 (0,0%)	0 (0,0%)	1 (100,0%)	
Расстройство настроения с психотическими симптомами, текущее, <i>n</i> (%)	0	39 (97,5%)	37 (100,0%)	25 (100,0%)	8 (100,0%)	109 (100,0%)	Chi2=1,766, <i>p</i> =0,622, <i>V</i> =0,127
	1	1 (2,5%)	0 (0,0%)	0 (0,0%)	0 (0,0%)	1 (100,0%)	
Нервная анорексия, текущая (за последние 3 месяца), <i>n</i> (%)	0	8 (20,0%)	9 (24,3%)	8 (32,0%)	1 (12,5%)	26 (100,0%)	Chi2=1,821, <i>p</i> =0,610, <i>V</i> =0,129
	1	32 (80,0%)	28 (75,7%)	17 (68,0%)	7 (87,5%)	84 (100,0%)	
Нервная булимия, текущая (за последние 3 месяца), <i>n</i> (%)	0	40 (100,0%)	37 (100,0%)	25 (100,0%)	8 (100,0%)	110 (100,0%)	Chi2=0,000, <i>p</i> =1,000, <i>V</i> =0,000
Нервная анорексия, тип с перееданием-очищением, текущая, <i>n</i> (%)	0	29 (72,5%)	26 (70,3%)	11 (44,0%)	7 (87,5%)	73 (100,0%)	Chi2=8,130, <i>p</i> =0,043, <i>V</i> =0,272
	1	11 (27,5%)	11 (29,7%)	14 (56,0%)	1 (12,5%)	37 (100,0%)	
Генерализованное тревожное расстройство, текущее (за последние 6 месяцев), <i>n</i> (%)	0	27 (67,5%)	24 (64,9%)	16 (64,0%)	4 (50,0%)	71 (100,0%)	Chi2=0,897, <i>p</i> =0,826, <i>V</i> =0,090
	1	13 (32,5%)	13 (35,1%)	9 (36,0%)	4 (50,0%)	39 (100,0%)	
Наличие медицинских, органических лекарственных причин, <i>n</i> (%)	0	39 (97,5%)	35 (94,6%)	22 (88,0%)	8 (100,0%)	104 (100,0%)	Chi2=3,216, <i>p</i> =0,360, <i>V</i> =0,171
	1	1 (2,5%)	2 (5,4%)	3 (12,0%)	0 (0,0%)	6 (100,0%)	

SCL-90-R							
Соматизация (Median [Q1; Q3])		0,71 [0,31; 1,08]	0,75 [0,42; 1,50]	1,17 [0,58; 1,67]	1,08 [0,83; 1,46]	0,83 [0,50; 1,40]	Kruskal: H=5,747, p=0,125, $\epsilon^2=0,053$
Обсессивно-компульсивное расстройство (навязчивости) (Median [Q1; Q3])		1,20 [0,67; 2,12]	1,40 [0,50; 1,90]	1,90 [1,30; 2,50]	1,45 [1,10; 2,05]	1,40 [0,70; 2,10]	Kruskal: H=4,714, p=0,194, $\epsilon^2=0,043$
Интерперсональная чувствительность (Median [Q1; Q3])		1,27 [0,64; 2,00]	1,11 [0,56; 2,00]	1,67 [1,00; 2,33]	1,00 [0,86; 1,19]	1,33 [0,59; 2,00]	Kruskal: H=3,422, p=0,331, $\epsilon^2=0,031$
Депрессия (Median [Q1; Q3])		1,38 [0,90; 1,87]	1,46 [0,62; 2,08]	1,85 [0,92; 2,54]	1,47 [1,25; 1,71]	1,50 [0,92; 2,13]	Kruskal: H=2,584, p=0,460, $\epsilon^2=0,024$
Тревожность (Median [Q1; Q3])		0,80 [0,35; 1,32]	0,70 [0,30; 1,70]	1,10 [0,50; 2,10]	1,30 [1,00; 1,30]	0,90 [0,40; 1,60]	Kruskal: H=3,326, p=0,344, $\epsilon^2=0,031$
Враждебность (Median [Q1; Q3])		0,75 [0,17; 1,17]	0,50 [0,17; 1,33]	0,67 [0,33; 1,67]	0,33 [0,17; 0,54]	0,67 [0,17; 1,33]	Kruskal: H=4,737, p=0,192, $\epsilon^2=0,043$
Навязчивые страхи (фобии) (Median [Q1; Q3])		0,43 [0,00; 1,00]	0,29 [0,00; 0,57]	0,43 [0,00; 1,00]	0,57 [0,25; 0,86]	0,43 [0,00; 0,86]	Kruskal: H=2,152, p=0,542, $\epsilon^2=0,020$
Параноидность (паранояльность) (Median [Q1; Q3])		0,42 [0,17; 1,17]	0,67 [0,17; 1,17]	1,17 [0,33; 1,50]	0,58 [0,46; 1,04]	0,67 [0,17; 1,17]	Kruskal: H=4,559, p=0,207, $\epsilon^2=0,042$
Психотизм (Median [Q1; Q3])		0,55 [0,40; 1,10]	0,50 [0,30; 1,40]	0,80 [0,40; 1,50]	0,55 [0,15; 0,75]	0,60 [0,30; 1,27]	Kruskal: H=2,142, p=0,544, $\epsilon^2=0,020$
Общий индекс тяжести симптомов GSI (Median [Q1; Q3])		0,95 [0,61; 1,50]	1,02 [0,41; 1,37]	1,27 [0,96; 1,94]	1,09 [0,83; 1,15]	1,02 [0,61; 1,58]	Kruskal: H=3,832, p=0,280, $\epsilon^2=0,035$
Общее число утвердительных ответов PST (Median [Q1; Q3])		46,50 [31,00; 60,25]	44,00 [22,00; 61,00]	62,00 [45,00; 73,00]	57,50 [36,50; 61,25]	50,00 [31,00; 64,00]	Kruskal: H=5,744, p=0,125, $\epsilon^2=0,053$
Индекс наличного симптоматического дистресса PSDI (Median [Q1; Q3])		1,96 [1,49; 2,28]	1,91 [1,50; 2,30]	2,12 [1,48; 2,42]	1,69 [1,59; 1,96]	1,92 [1,49; 2,30]	Kruskal: H=1,329, p=0,722, $\epsilon^2=0,012$

Примечание: M.I.N.I. — The Mini-International Neuropsychiatric Interview; SCL-90-R — Symptom Check List-90-Revised; ИМТ — индекс массы тела; РПП — расстройство пищевого поведения; ЭПС — экстрапирамидные симптомы.